

Weak Gravitational Field in Finsler - Randers Space and Raychaudhuri Equation

P. Stavrinou*

Abstract

The linearized form of the metric of a Finsler - Randers space is studied in relation to the equations of motion, the deviation of geodesics and the generalized Raychaudhuri equation are given for a weak gravitational field. This equation is also derived in the framework of a tangent bundle.

By using Cartan or Berwald-like connections we get some types "gravito - electromagnetic" curvature. In addition we investigate the conditions under which a definite Lagrangian in a Randers space leads to Einstein field equations under the presence of electromagnetic field.

Finally, some applications of the weak field in a generalized Finsler spacetime for gravitational waves are given.

keywords: Finsler geometry, weak gravitational field, Raychadhuri equation.

*Department of Mathematics, University of Athens, 15784 Panepistimiopolis, Athens, Greece. Email: pstavrin@math.uoa.gr

1 Introduction

In the framework of general relativity, weak fields and gravitational waves have been studied by many authors decades ago e.g.[1],[2],[3]. One of the fundamental problems in general relativity is the study of gravitational waves. The existence of gravitational waves in linear versions of the theory was already known in the early days of general relativity. First Einstein considered in a Minkowski spacetime with a metric $n_{\mu\nu}$ a small perturbation $\epsilon_{\mu\nu}$ such that the induced field $\alpha_{\mu\nu} = n_{\mu\nu} + \epsilon_{\mu\nu}$ with $|\epsilon_{\mu\nu}| \ll 1$ obeys by perturbation in linearized equations of motion. The linearized theory of gravity is an important theory because it can be utilized as a foundation for “deriving” General Relativity. By using the linearized field theory of gravitation some observable phenomena of our solar system and the universe can be detected [4],[5] The weak field limit at a Finsler space-time has been studied in [6],[7] and in the tangent bundle of a Finsler space by [8],[9], [10].

In our study we consider a pseudo- Finsler - Randers space-time of metric function [11]

$$\mathcal{F}(x, y) = \sqrt{\alpha_{ij}(x)y^i y^j} + kA_i(x)y^i \quad (1)$$

in the weak field limit where α_{ij} represents pseudo-Riemannian metric, $y^i = \frac{dx^i}{d\lambda}$ (in applications y^i represent velocity), λ a parameter along the curve and k a constant, A_i represents the electromagnetic potential which is connected with the electromagnetic field by $F_{ij} = \frac{\partial A_j}{\partial x^i} - \frac{\partial A_i}{\partial x^j}$.

A Finsler-Randers space (FR) constitutes an important category of Finsler spaces from mathematical and physical perspective e.g.[12],[13],[14],[15]. In a FR space the condition of symmetry for the fundamental function $\mathcal{F}(x, y)$ is not satisfied $\mathcal{F}(x, y) \neq \mathcal{F}(x, -y)$. Causal considerations in pseudo-Finsler space-time that include only symmetries of fundamental functions [16] are very restricted and exclude by studying pseudo-FR spaces.[12],[15],[17].

FR space can play a significant role in the theory of weak field and gravitational waves since ”a gravito - electromagnetic field” is intrinsically included in its metric. Einstein’s General Relativity shows indeed that gravito - magnetic field may be associated with mass currents [18],[19]. As well a gravito - magnetic force was postulated as an explanation for the anomalous precession of Mercury’s perihelion [20]. In addition, previous works [21], [22], [23] ,[24], [25] showed how the electric and magnetic parts of the curvature tensors were related to the electric and magnetic parts of the gravitational field as well as with gravitatalional waves.[26]

This paper is organized as follows: we deal with the linearized metric form of a pseudo FR spacetime in relation to the deviation of geodesics and Raychaudhuri equation. In this approach we extend a previous consideration which was given in [6]. By using Cartan and Berwald -like connections we get “gravito - electromagnetic curvatures” for this space. We derive the equations of motion and the Raychaudhuri equation in the framework of a tangent bundle of a n -dimensional manifold M . We also give the linearized connection coefficients as well as establishing the Lorentz equation of the weak field. In addition we attribute some physical interpretations in the geometrical concepts under consideration.

Moreover in the framework of the weak gravitational field of a generalized Finsler spacetime some applications for gravitational waves are given.

2 Linearized field theory of Randers space-time

The behavior of particles in a gravitational and electromagnetic field is expected to indicate that the physical geometry in the direction of a geometrical unification is the Finsler geometry.

In a Finsler space the metric function $\mathcal{F}(x, y)$ can be considered as a potential function since the metric tensor (gravitational potential)

$$g_{ij}(x, y) = \frac{1}{2} \frac{\partial^2 \mathcal{F}^2}{\partial y^i \partial y^j} \quad (2)$$

is produced by this function. The metric of a Randers space is given by virtue of (2) in the form [12],[27].

$$g_{ij} = \alpha_{ij} + \frac{2k}{\sigma} y^s \alpha_{s(i} A_{j)} + k^2 A_i A_j + \frac{k}{\sigma} y^l A_l m_{ij} \quad (3)$$

where $\sigma = \sqrt{\alpha_{ij} y^i y^j}$, $m_{ij} = \alpha_{ij} - \sigma^{-2} \alpha_{is} \alpha_{jl} y^s y^l$ and $A_{(ij)} = \frac{1}{2} (A_{ij} + A_{ji})$. We observe that the presence of an electromagnetic field in a region of spacetime breaks the isotropy and the description of spacetime is given by two metrics, one of which has a pseudo-Riemann structure $a_{ij}(x)$ that corresponds to a motion of a particle with mass m in the gravitational field and the second is metric of a charged particle of mass m that corresponds to a Finsler space of

metric $g_{ij}(x, v)$ which represents a dynamical field. Connection coefficients of pseudo FR space are produced by those metrics. In a FR space the second term $y^i b_i$ can represent a measure of a cosmological anisotropy, a magnetic field or a spin-velocity. This consideration is analogous to Rosen's (1940) in which at each point of space-time a Euclidean and pseudo-Riemannian metric corresponds in each point of space-time. (Bimetric theory)

Finsler spaces are endowed with Cartan, Berwald connections and other different types of connections. Cartan connection has very important properties (metric compatibility) for models are closely related to standard physics [28],[29]. Berwald connection is not generally compatible with the metric structure on total space, since it has "weak" compatibility only on the h-space on the tangent bundle. However for the case of a standard model extension a Berwald structure can be used for Lorentz violation in relation to gravitational waves e.g. [30]

The explicit form of Randers connection of Berwald-like coefficients can be given in the form [31]

$$L_{ij}^l = \alpha_{ij}^l + E_{ij}^l, \quad (4)$$

where α_{ij}^l are the Riemannian Cristoffel symbols and E_{ij}^l are given by

$$E_{ij}^l = \frac{1}{2} (\alpha_{ij} y^k F_k^l + u_i F_j^l + u_j F_i^l) \alpha^{-1} - \frac{1}{2} u_i u_j y^k F_k^l \alpha^{-3} \quad (5)$$

with $u_i = u_i(x, y) = \frac{\alpha_{ij} y^j}{\alpha(x, y)}$ $\alpha = \alpha(x, y) = (\alpha_{ij}(x) y^i y^j)^{1/2}$. We note from (4) and (5) that the electromagnetic field enters in the connection coefficient of this space.

The geodesics of the Randers space are produced by the first variation of the action corresponding to the Lagrangian.

$$\frac{d}{d\lambda} \left(\frac{\partial \mathcal{F}}{\partial y^m} \right) - \frac{\partial \mathcal{F}}{\partial x^m} = 0 \quad (6)$$

$$\frac{dy^m}{d\lambda} + L_{ij}^m(x, y) y^i y^j = 0 \quad (7)$$

Because of (7) we get the well known Lorentz equation

$$\frac{d^2 x^m}{d\lambda^2} + \alpha_{ij}^m \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda} + k F_j^m \frac{dx^j}{d\lambda} = 0 \quad (8)$$

Eq.(8) represents the equation of motion of a charged particle in a gravitational and electromagnetic field, where λ represents an affine parameter.

The equations of motion (8) under the perturbations h_{ij} can be written in the form

$$\frac{d^2 x^m}{d\lambda^2} + \tilde{\alpha}_{ij}^m \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda} + k F_j^m \frac{dx^j}{d\lambda} = 0 \quad (9)$$

where $\tilde{\alpha}_{ij}^m = \frac{1}{2} n^{ml} (\partial_i h_{jl} + \partial_j h_{im} - \partial_l h_{ij})$ are the connection coefficients of the weak metric $\tilde{\alpha}_{ij}$. The Eq. (9) is useful for studying gravitational waves in a pseudo-FR space.

The curvature tensor of a Randers space can be produced by using Berwald-like connection coefficients in analogous to the form [31] for a weak field

$$H_{hjk}^i = R_{hjk}^i + E_{hjk}^i \quad (10)$$

where R_{hjk}^i is the Riemannian curvature tensor and E_{hjk}^i is given by

$$\begin{aligned} E_{hjk}^i &= \frac{1}{2} Q_{[jk]} (F_h^i F_{jk} + \alpha_{hk} F_j^m F_m^i - F_{hj} F_k^i) \\ &+ Q_{[jk]} [(u_h \nabla_k F_j^i + y^m \alpha_{hj} \nabla_k F_m^i + u_j \nabla_k F_h^i) \alpha^{-1} - y^m u_h u_j \alpha^{-3} \nabla_k F_m^i] \end{aligned} \quad (11)$$

with $\alpha = \alpha(x, y) = (\alpha_{ij} y^i y^j)^{1/2}$.

Relation (10) is rewritten

$$H_{hjk}^i(x, y) = \Lambda_{hjk}^i(x) + Q_{[jk]} (u_h \nabla_k F_j^i + \dots - \nabla_k F_m^i) \quad (12)$$

with $\Lambda_{hjk}^i = R_{hjk}^i + \frac{1}{2} Q_{[jk]} (F_h^i F_{jk} + \dots - F_{hj} F_k^i)$ and $Q_{[jk]} = \frac{1}{2} (Q_{kj} - Q_{jk})$. Applying the condition

$$Q_{[jk]} (u_h \nabla_c F_j^i + \dot{x}^m \tilde{\alpha}_{hj} \nabla_k F_m^i + u_j \nabla_k F_h^i) \tilde{\alpha}^{-1} - \dot{x}^m u_h u_j \tilde{\alpha}^{-3} \nabla_k F_m^i = 0 \quad (13)$$

in (10) we get the Lagrangian of the classical gravitational and electromagnetic fields

$$\Lambda = \Lambda_{hji}^i \alpha^{hj} = R + k F_{mn} F^{mn}, \quad k \text{ constant.} \quad (14)$$

The variation of action's integral

$$\delta I = \delta \int \Lambda \sqrt{|\tilde{g}|} d^4 x \quad (15)$$

$$g = \det(a_{ij}) \quad (16)$$

leads to the weak field equations

$$\left(R_{mn} - \frac{1}{2} \tilde{\alpha}_{mn} R \right) + k \left(F_{nr} F_m^r - \frac{1}{4} \tilde{\alpha}_{mn} F_{rs} F^{rs} \right) = 0 \quad (17)$$

which are Einstein - Maxwell field equations of the “gravito - electromagnetic” of the Randers space for the vacuum under the condition (13). These equations have the same form as in the Riemannian ansatz in the presence of an electromagnetic field.

From a physical point of view the curvature H_{hjk}^i can be considered as a “gravito - electromagnetic curvature” of the space. We can say that it involves a gravito - electromagnetic “current” source. The metric of a weak gravitational field can be decomposed into the flat Minkowski metric plus a small perturbation

$$\tilde{\alpha}_{ij} = n_{ij} + h_{ij}, \quad |h_{ij}| \ll 1. \quad (18)$$

Under a linearized approach of the gravitational field, the Randers metric function can be written in the form of a first approximation of the Riemannian metric α_{ij}

$$\mathcal{F}(x, y) = \sqrt{(n_{ij} + h_{ij}(x, v)) v^i v^j} + k A_i v^i \quad (19)$$

where $v^i = dx^i/d\tau$ is the 4-velocity of the particle, $n_{ij} = \text{diag}(1, -1, -1, -1)$ is the Minkowski metric, $|h_{ij}| \ll 1$ represents small perturbations to the flat spacetime metric and k is a constant. The linearized form of the metric tensor, as introduced by (3), becomes

$$g_{ij} = n_{ij} + h_{ij} + \frac{2k}{\sigma} v^s n_{s(i} A_{j)} + k^2 A_i A_j + \frac{k}{\sigma} v^l A_l \theta_{ij} \quad (20)$$

where $\sigma = \sqrt{n_{ij} v^i v^j}$, $\theta_{ij} = n_{ij} - \sigma^{-2} n_{si} n_{jl} v^s v^l$ and $a_{(ij)} = \frac{1}{2} (a_{ij} + a_{ji})$.

Considering in (20) the case where $v^s = (1, 0, 0, 0)$ we get the Finslerian potential g_{00} of the Randers space for a test material point in the static case

$$g_{00} = 1 + h_{00} + k^2 \phi^2 + 2k\phi \quad (21)$$

with $\phi = A_0$. In the case of $\phi = 0$ we get $g_{00} = 1 + h_{00}$. This relation is useful in order to derive the Riemannian or Newtonian limit from the equation of motion in a Randers space. In this case the equation of motion has the form

$$\dot{v}^l + L_{00}^l v^0 v^0 = \dot{v}^l + L_{00}^l = 0 \quad (22)$$

with

$$L_{00}^\mu = -\frac{1}{2}n^{\mu\lambda}\partial_\lambda h_{00}. \quad (23)$$

The full interpretation of L_{ij}^l is given by (4).

The Finslerian potential g_{00} takes the value 1 for the values of electromagnetic vector potential

$$\phi_{1,2} = \left(-1 \pm (1 - h_{00})^{1/2}\right) (k)^{-1}. \quad (24)$$

The Cristoffel symbols and the curvature tensor of the linearized Randers space will take the following form. By using (4) and (10) we get

$$\tilde{L}_{lj}^i = \tilde{a}_{lj}^i + h_{lj}^i \quad (25)$$

$$\tilde{H}_{ljk}^i = \tilde{R}_{ljk}^i + h_{ljk}^i \quad (26)$$

where

$$\tilde{R}_{ljk}^i = \frac{1}{2}n^{is} (\partial_{[kl}^2 h_{sj]} - \partial_{[js}^2 h_{lk]}) \quad (27)$$

\tilde{a}_{jk}^i are the linearized Riemannian Christoffel symbols and the curvature tensor. From (25) and (26) the rest terms will be given in the form

$$h_{ij}^m = \frac{1}{2} (n_{ij}v^k F_k^m + u_i F_j^m + u_j F_i^m) n^{-1} - \frac{1}{2} u_i u_j v^k F_k^m n^{-3} \quad (28)$$

$$\begin{aligned} h_{hjk}^i &= \frac{1}{2} Q_{[jk]} (F_h^i F_{jk} + n_{hk} F_j^m F_m^i - F_{hj} F_k^i) \\ &+ Q_{[jk]} (u_h \partial_k F_j^i + v^m n_{hj} \partial_k F_m^i + u_j \partial_k F_h^i) n^{-1} - v^m u_h u_j n^{-3} \partial_k F_m^i \end{aligned} \quad (29)$$

By using Cartan covariant differentiation in a Randers space we can express the third curvature tensor of Cartan \bar{R}_{jkl}^i in the form [32]

$$\begin{aligned} \bar{R}_{jhk}^i &= R_{jhk}^i + D_{jh|k}^i - D_{jk|h}^i + D_{hk}^i D_{jk}^m - D_{mk}^i D_{jh}^m \\ &+ C_{jm}^i (R_{0hk}^m + D_{0h|k}^m - D_{0k|h}^m + D_{sh}^m D_{0k}^s - D_{sk}^m D_{0h}^s) \end{aligned} \quad (30)$$

where R_{jhk}^i is the Riemannian curvature, the symbol $|$ represents the Cartan covariant derivative and D_{jk}^i is the difference tensor of the Finslerian

gravitational field given by

$$\begin{aligned}
D_{jk}^i = & k\ell^i A_{(j,k)} + \frac{1}{2}k (\omega_j^i A_{0,k} + \omega_k^i A_{0,j} - \omega_{jk} A_{0,s} g^{is}) - C_{jk}^i \\
& + g^{is} k (A_{[s,j]} \ell_k + A_{[s,k]} \ell_j) + \frac{k}{\tau} \alpha^{mt} [g^{is} (A_{([t,s])} + A_{[t,0]} p_s \tau) C_{jkm} \\
& - C_{km}^i (A_{[t,j]} + A_{[t,0]} p_j \tau) - C_{jm}^i (A_{[t,k]} + A_{[t,0]} p_k \tau)]
\end{aligned} \tag{31}$$

where $\omega_{ij}(x, y) = \tau (a_{ij} - k^2 A_i A_j)$, $\tau = \mathcal{F}/\alpha^{1/2}$ and $\ell^i = y^i/\mathcal{F}$. In a linearised form the Cartan curvature tensor is expressed by

$$\tilde{K}_{jhk}^i = \tilde{R}_{jhk}^i + \tilde{D}_{jh|k}^i - \tilde{D}_{jk|h}^i + \tilde{C}_{jm}^i \left(\tilde{R}_{0hk}^m + \tilde{D}_{0h|k}^m - \tilde{D}_{0k|h}^m \right) \tag{32}$$

where \tilde{R}_{jhk}^i is given by (27) and $\tilde{D}_{jh}^i, \tilde{C}_{jm}^i$ represent the weak difference tensor and the weak Cartan connection coefficients, $\tilde{C}_{ijk} = \frac{1}{2} \frac{\partial \tilde{g}_{ij}}{\partial y_k}$.

In (30) we have ignored terms $\tilde{D} \cdot \tilde{D}$ because of the condition $|h_{ij}| \ll 1$. The Ricci tensor of the weak "gravito - electromagnetic" field is given by

$$\tilde{K}_{ij} = \tilde{K}_{ijs}^s. \tag{33}$$

For a perfect fluid moving in a Randers space with Cartan curvature K_{jhk}^i Einstein's equations can be given in the form of weak field

$$\tilde{K}_{il} = k \left(T_{il}(x, V(x)) - \frac{1}{2} T_{\kappa}^{\kappa} g_{ij} \right) \tag{34}$$

with $\tilde{K}_{il} = \tilde{R}_{il} + E_{il}$, where \tilde{K}_{il} is the Ricci tensor, \tilde{R}_{il} is the Riemannian one, E_{il} is the contraction of E_{jkl}^i by (10) and T_{il} the energy - momentum tensor of FR space. Randers type space-time in cosmological considerations for a weak anisotropic field u^a with $\|u^a\| \ll 1$ has been studied in [13].

3 Weak Deviation of geodesics. Raychaudhuri equation

The deviation of geodesics play an important role in General Relativity and Gravitation. In the Finslerian space-time it has been studied from

mathematical and physical point of view [33],[34],[35],[36]. In a pseudo-Randers space the deviation of geodesics can be expressed by using "gravito-electromagnetic" curvature (10)

$$\frac{\delta^2 \xi^i}{\delta \lambda^2} + \bar{H}_{j^h k}^i(x, v) \xi^j v^h v^k = 0 \quad (35)$$

where $\frac{\delta \xi^i}{\delta \lambda} = \xi_{|h}^i v^h$ and $\xi_{|h}^i = \frac{\partial \xi^i}{\partial x^h} + \bar{R}_{hk}^i(x, v) \xi^k$, with λ affine parameter. \bar{R}_{hk}^i are the Cartan connection coefficients, ξ^i represents the deviation vector and v^k the tangent vectors of a geodesic surface included in the Randers spacetime. We note from (35) that the deviation equation has two terms. The first term corresponds to the gravitational deviation, that will be observed if there is no electromagnetic field and it is associated with the R_{hjk}^i part of the curvature tensor. The other term corresponds to a mixed geometrical and electromagnetic deviation and is associated with the E_{hjk}^i part of the curvature tensor. The second term of deviation is connected to the force that two freely falling charged particles would exert to each other. Studying this case in a Riemannian spacetime has the consequence that the force does not necessarily result as a natural geometric effect as it does in a Finsler spacetime. If the E_{hjk}^i vanishes then the deviation equation is reduced to the well known one of the Riemannian spacetime, namely

$$\frac{\delta^2 z^i}{\delta u^2} + R_{jkl}^i z^j v^k v^l = 0 \quad (36)$$

Physically this means that we have two freely falling particles in the tidal field R_{hjk}^i of spacetime. In the case where R_{hjk}^i vanishes, we infer that the first term of the Randers metric corresponds to a Minkowski metric and the Finsler - Randers space becomes v - locally Minkowski [37].

$$\mathcal{F}(x, v) = \sqrt{n_{\mu\nu} v^\mu v^\nu} + k A_i(x) v^i \quad (37)$$

The only force that influences the two charged particles is due to the presence of the charged electromagnetic field. The deviation equation takes the form

$$\frac{\delta^2 z^i}{\delta u^2} + E_{jkl}^i z^j v^k v^l = 0 \quad (38)$$

where E_{jkl}^i is given by (11). In this case the geometrical properties of the field are characterized by a homogeneous and anisotropic space. The metric fundamental tensor depends only on the velocities, which produce the

anisotropic properties of the curved Finsler spacetime. Consequently there exists a frame of reference, where \bar{R}_{hjk}^i vanishes. Under these circumstances the geodesic coordinates can be introduced for particles moving along these geodesics.

In a cosmological consideration the formula(37) can be given by $\mathcal{F}(x, V) = \sqrt{n_{\mu\nu}V^\mu V^\nu} + kW_i(x)V^i$ where $V = \tilde{H}d$ represents cosmological velocity depending on the cosmological Hubble parameter \tilde{H} which is defined in the anisotropic Randers space-time with $\tilde{H} = \sqrt{H^2 + Hz_t}$ cf. [13], W_i represents an anisotropic field, Z_t the variation of anisotropy and d the distance. Such a consideration can be provided by a Finslerian osculating geometrical framework.If this field W_i comes from by a curl the geodesics of this model are Riemannian. Gravitational waves in locally anisotropic spaces generate polarization patterns of the cosmic microwave background.

It is well known that the gravitational waves are connected to the deviation of geodesics. In order to study the weak field limit of a Randers space related to the deviation of the charged particles it is necessary to take into account relations (26) and (35). This is reasonable since in order to detect a gravitational wave at least two particles are needed.

Thus the deviation of geodesics of the weak Randers space is written in the form

$$\frac{d^2 z^i}{d\tau^2} + \tilde{h}_{ijm}^i \frac{dz^j}{d\tau} \frac{dx^l}{d\tau} \frac{dx^m}{d\tau} = 0 \quad (39)$$

$$\frac{d^2 z^i}{d\tau^2} + (\epsilon_{ijm}^i + h_{ijm}^i) \frac{dx^l}{d\tau} \frac{dz^j}{d\tau} \frac{dx^m}{d\tau} = 0 \quad (40)$$

If we consider our test particles to be moving slowly then we can express the 4-velocities as a unit vector in the time direction. Hence we write

$$\frac{dx^i}{d\tau} = (1, 0, 0, 0). \quad (41)$$

In order to compute the Riemannian tensor in a first approximation we get from (40)

$$\epsilon_{0j0}^i = \frac{1}{2}n^{ik} (\epsilon_{jk,00} - \epsilon_{0j,k0} - \epsilon_{0k,0j} + \epsilon_{00,kj}) \quad (42)$$

In our case $\epsilon_{i0} = 0$ and the Riemannian tensor takes the form

$$\epsilon_{0j0}^i = \frac{1}{2}\epsilon_{j,00}^i. \quad (43)$$

The second term of (40) $h_{l_j m}^i$ because of (41) and (29) becomes

$$h_{0j0}^i = 2F_0^i F_j^0 + u_j \partial_0 F_0^i. \quad (44)$$

Furthermore (40) will take the form because of (43) and (44)

$$\frac{\partial^2 z^i}{\partial \tau^2} + \left(\frac{1}{2} \frac{\partial^2 \epsilon_j^i}{\partial \tau^2} + 2F_0^i F_j^0 + u_j \frac{\partial F_0^i}{\partial \tau} \right) \frac{\partial z^j}{\partial \tau} = 0. \quad (45)$$

The equation (45) coincides with the corresponding equation for a weak field limit of the Riemannian case, which is given in its full form by

$$\frac{d^2 n^\mu}{ds^2} + R_{\nu\kappa\lambda}^\mu n^\kappa v^\nu v^\lambda = \Phi^\mu \quad (46)$$

with

$$\Phi^\mu = k \left(\frac{dF_\kappa^\mu}{ds} v^\kappa + F_\nu^\mu F_\kappa^\nu v^\kappa \right), \quad k : \text{constant}. \quad (47)$$

In general Φ^μ represents a non-gravitational force, for instance a spring. The difference between (46) and (35) is that in (46) the electromagnetic field has been added *ad hoc*. This means that the term Φ^μ plays the role of the interaction external force between two nearby charged masses, moving in non-geodesical paths. In the equation (35) of the Randers space the electromagnetic field is incorporated in the geometry. In this approach the two charged masses move in geodesics of the Finsler space and their relative acceleration is determined by the curvature of the gravitational and electromagnetic fields, which is produced by the energy momentum tensor. Randers-type spaces best express a profound relation between physics and geometry.

An extension of the geodesic deviation equations constitutes the Raychadhuri equation. Raychadhuri equation is of important significance in Relativity theory and Cosmology because of its connection with singularities e.g. [38],[39],[40]. In Finsler-Randers space-time this equation has been studied in a previous paper [43]. Its form in the weak Finslerian limit is given by

$$\frac{d\tilde{\theta}}{d\tau} = -\frac{1}{3}\tilde{\theta}^2 - \tilde{\sigma}_{ik}\tilde{\sigma}^{ik} + \tilde{\omega}_{ik}\tilde{\omega}^{ik} - K_{i\ell}V^iV^\ell + \dot{V}_{;i}^i \quad (48)$$

where $K_{i\ell}$ represents the weak Cartan tensor, $\tilde{\theta}$, $\tilde{\omega}_{ik}$, $\tilde{\sigma}_{ik}$ are the expansion, vorticity

and the shear are defined by the following forms:

$$\tilde{\theta} = \Lambda_{ij} h^{ij} = V_{|i}^i - C_{im}^i \dot{V}^m \quad (49a)$$

$$\tilde{\omega}_{ik} = \Lambda_{[ik]} + \dot{V}_i V_k - \dot{V}_k V_i \quad (49b)$$

$$\tilde{\sigma}_{ik} = \Lambda_{(ik)} - \frac{1}{3} \tilde{\theta} h_{ik} - 2C_{ikm} V^m - \dot{V}_i V_k - \dot{V}_k V_i \quad (49c)$$

where $\Lambda_{(ik)} = V_{i;k}$ is the covariant derivative of the osculating Riemannian space V^i a unit vector $V^i V_i = 1$. In the case of geodesics the last term $\dot{V}_{;i}^i$ vanishes in (48) and (49). The introduction of Cartan tensor in (48) assigns an anisotropic structure for the Raychaudhuri equation. The linearized Raychaudhuri equation in a Randers spacetime in a first approach is expressed without vorticity and shear by

$$\frac{d\tilde{\theta}}{d\tau} = -\frac{1}{3} \tilde{\theta}^2 - \tilde{K}_{il} V^i V^l \quad (50)$$

In the case that $\dot{\tilde{\theta}} = 0, \tilde{\sigma}_{ij} = 0, \tilde{\omega}_{ij} = \text{constant}$, from (48) the tidal field $K_{il} V^i V^l$ is due to the vorticity $\tilde{\omega}$ which plays the role of vacuum energy (cosmological constant). It is analogous to a centrifugal field of the Newtonian theory. It counterbalances the tidal field.

Remark The fundamental sense of photon surfaces and their geometry has been defined and developed in [41],[42] for a timelike surface in a spherical symmetric space with determined properties.

In a pseudo-Finsler space-time M with spherically symmetric metric [35] in which $\tilde{\sigma}_{ij} = 0$ and $\tilde{\omega}_{ij} = 0$, we can analogously consider a Finslerian photon surface S , where S represents a timelike surface of M . Here the Raychaudhuri equation takes the form

$$\frac{d\theta_{(2)}}{d\tau} = -\frac{1}{2} \theta_{(2)}^2 - K_{il}^{(3)} X^i X^l, \quad (51)$$

where $\theta_{(2)}$ denotes the expansion of a vector field X in the surface S , τ the affine parameter and $K_{il}^{(3)}$ the Ricci tensor of Cartan curvature.

On a physical viewpoint the anisotropic Cartan tensor is introduced in the geometry of spacetime because of a primordial vector field in the Finsler-Randers spacetime. Such a case has been studied in [13] where the linearized

Raychaudhuri equation stands

$$\frac{1}{3}\tilde{\theta}^2\tilde{q} = 4\pi G(\mu + 3P)\frac{H}{\tilde{H}} + f(\alpha, \dot{\alpha}, \ddot{\alpha}, z_t) \quad (52)$$

where $\tilde{q}, \mu, p, \alpha, z_t$ represent the deceleration parameter, the density of matter, the pressure, the scale factor and the variation of anisotropy, which is connected to Cartan connection component $z_t = C_{000,0}$.

The Raychaudhuri equations can also be derived in the framework of a tangent bundle TM of a n -dimensional manifold by using of d-connection and the Ricci-identities. In this case we consider the d-curvature R_{jkl}^i and the Ricci identities for a tangent horizontal vector field $X = X^H = X^i\delta/\delta x^i$ along a congruence of geodesics on TM [37]. So we have the relation

$$X_{|kl}^i - X_{|lk}^i = R_{jkl}^i X^j - T_{kl}^h X_{|h}^i - R_{kl}^a X^i|_a, \quad (53)$$

or

$$X^l X_{|kl}^i = X_{|lk}^i X^l + R_{jkl}^i X^j X^l - T_{kl}^h X_{|h}^i X^l - R_{kl}^a X^i|_a X^l. \quad (54)$$

Because of geodesics the relation $(X^l X_{|l}^i)|_k = 0$ is valid so we have

$$X^l X_{|kl}^i = -X^l X_{|k} X_{|l}^i + R_{jkl}^i X^j X^l - T_{kl}^h X_{|h}^i X^l - R_{kl}^a X^i|_a X^l. \quad (55)$$

Taking the trace of the previous equation we have

$$X^l \tilde{h}_i^k X_{|kl}^i = -X_{|k}^l X_{|l}^i \tilde{h}_i^k + R_{jkl}^i X^j X^l \tilde{h}_i^k - T_{kl}^h X_{|h}^i X^l \tilde{h}_i^k - R_{kl}^a X^i|_a X^l \tilde{h}_i^k. \quad (56)$$

We decompose the h -covariant derivative with kinematical terms

$$X_{|l}^i = \frac{1}{n-1}\tilde{\Theta}\tilde{h}_l^i + \tilde{\sigma}_l^i + \tilde{\omega}_l^i, \quad (57)$$

where $\tilde{\Theta}, \tilde{\sigma}, \tilde{\omega}$ represent the expansion, shear and vorticity for the extended congruence of geodesics on TM, which are defined as

$$\tilde{\Theta} = X_{|l}^i \tilde{h}_i^l, \quad (58)$$

$$\tilde{\sigma}_{il} = X_{i|l} + X_{l|i} - \frac{1}{3}\tilde{\Theta}\tilde{h}_{il}, \quad (59)$$

$$\tilde{\omega}_{il} = X_{i|l} - X_{l|i}, \quad (60)$$

with

$$\tilde{h}_{il} = g_{il} - X_i X_l \quad (61)$$

the projection operator, h_{il} gives us $\tilde{h}_{il}X^i = 0$ for a normalized X_i . In the above mentioned relations, (57), (58), we used

$$\tilde{h}_l^i = g^{ik}\tilde{h}_{lk}. \quad (62)$$

We finally get from (56)

$$\begin{aligned} X^l\tilde{\Theta}|_l &= \frac{d\tilde{\Theta}}{d\tau} = R_{kl}X^kX^l - T_{li}^hX^i|_hX^l - R_{li}^aX^i|_a - X^l|_kX^k|_l = R_{kl}X^iX^l \\ &- T_{li}^h\left(\frac{1}{n-1}\tilde{\Theta}\tilde{h}_h^i + \tilde{\sigma}_h^i + \tilde{\omega}_h^i\right) - R_{li}^a\left(\frac{1}{n-1}\Theta\tilde{h}_a^i + \sigma_a^i + \omega_a^i\right) - \frac{1}{n-1}\tilde{\Theta}^l - \sigma_k^l\sigma_l^k - \omega_k^l\omega_l^k. \end{aligned} \quad (63)$$

where we put

$$X^i|_a = \frac{1}{n-1}\tilde{\Theta}\tilde{h}_a^i + \tilde{\sigma}_a^i + \tilde{\omega}_a^i,$$

with $\tilde{h}_a^i = h_a^b\delta_c^i\delta_b^c$, $\sigma_a^i = \sigma_a^b\delta_c^i\delta_b^c$, $\omega_a^i = \omega_a^b\delta_c^i\delta_b^c$ and δ_c^i represent the generalized Kronecker symbols connecting with h-bases and v-bases. The equation (63) is the Raychadhuri equation for the horizontal space of the tangent bundle.

$$R_{li}^a = \frac{\partial N_l^a}{\partial x^i} - \frac{\partial N_i^a}{\partial x^l} \quad (64)$$

represents the curvature of non-linear connection.

By using of d-curvature S_{bcd}^a , the Ricci identities are written

$$X^a|_{bc} - X^a|_{cb} = S_{abc}^dX^d - S_{bc}^dX^a|_d, \quad (65)$$

where the vector field $X = X^a\frac{\partial}{\partial y^a}$ belongs to the vertical space $S_{bc}^a = C_{bc}^a - C_c^ab$ represent the torsion and C_{bc}^a the d-connection coefficients of the vertical space. In analogy to the consideration of Finslerian fluids cf.[43] we can get the Raychaudhuri equations. For the definitions of the decomposition of vertical covariant derivative of vertical geodesics expansion, shear and rotation we use the relations

$$X^a|_b = \frac{1}{3}\Theta h_b^a + \sigma_b^a + \omega_b^a \quad (66)$$

where the expansion Θ is given by

$$\Theta = X^a|_bh_a^b = X^a|_a, \quad (67)$$

$h_{ab}(x, y)$ represents the v-metric on TM which is connected with the h-metric $g_{ij}(x, y) = \delta_i^a \delta_j^b h_{ab}(x, y) \sigma_{ab}$ and ω_{ab} . We define the shear and rotation by

$$\sigma_{ab} = X_a|_b + X_b|_a - \frac{1}{3}\Theta h_{ab}, \quad (68)$$

$$\omega_{ab} = X_a|_b - X_b|_a. \quad (69)$$

Because of the above mentioned relations one obtains an expression for the Raychaudhuri equation in the vertical space in the forms

$$\begin{aligned} X^c \Theta|_c &= \frac{d\Theta}{d\tau} = S_{dc} X^d X^c - S_{ca}^d X^a|_d X^c - X^c|_b X^b|_c \\ &= -\frac{1}{n-1} \Theta^2 - \sigma_b^a \sigma_a^b - \omega_b^a \omega_a^b + S_{dc} X^d X^c - S_{ca}^d X^c \left(\frac{1}{n-1} \Theta h_b^a + \sigma_b^a + \omega_b^a \right) \end{aligned} \quad (70)$$

The Raychaudhuri equation can also be derived on the tangent bundle of a Finsler-Randers space-time as well as for its weak field limit by considering the analogous curvature in the rel. (63).

Applications:

1. In the (x^0, x^1, x^2, x^3) coordinates of an inertial frame generalized Finsler metrics that are close to the flat metric can be written

$$g_{\alpha\beta}(x, y) = n_{\alpha\beta} + \epsilon_{\alpha\beta}(x, y) \quad (71)$$

where $y = \frac{dx}{dt}$ and $\epsilon_{\alpha\beta}(x, y)$ are small anisotropic perturbations to the flat spacetime. These metric perturbations describe a gravitational wave. The line - element for a plane gravitational wave spacetime can be expressed in the form

$$ds^2 = dt^2 - (1 + f[(x_0, x_3), y] dx_1^2) - (1 - f[(x_0, x_3), y] dx_2^2) + dx_3^2 \quad (72)$$

where the function $f(x, y) = \epsilon_{\alpha\beta}(x, y)$, with

$$|f[(x^0 - x^3), y]| \ll 1. \quad (73)$$

If the wave has a definite frequency ω , amplitude α and phase δ we can write

$$f[(x^0 - x^3), y] = \alpha \sin[\omega(x^0 - x^3) + \delta] y. \quad (74)$$

If we focus on the plane $z = 0$, because of the weak field (71) we can write

$$\alpha \sin(\omega t + \delta) y^k = h_{ij}(t, 0) n^i n^j y^k \quad (75)$$

where n^i is a unit vector on the tangent plane \mathbb{R}^2 , which is identical to the euclidean plane \mathbb{R}^2 . Equation (75) shows how gravitational waves can be detected by observing the relative motion of two test masses. This is one way of defining the local curvature of spacetime in general. Before the wave passes the spacetime is flat. After the wave passes the motion of a test mass is given by

$$\frac{dk^i}{d\tau} + \tilde{C}_{\alpha\beta}^i k^\alpha k^\beta = 0 \quad (76)$$

where $\tilde{C}_{\alpha\beta}^i$ are the linearized Cartan connection coefficients, which denote a propagation of gravitational waves in an anisotropic spacetime.

2. In the previous application the coordinates x^i , $i = 0, 1, 2, 3$ are independent of time. Distances between test masses in Euclidean plane can be calculated from $X(t), Y(t)$ coordinates by using a Randers metric. We can put the coordinates in the form

$$X = (1 + \frac{1}{2}\alpha \sin \omega t) x^1 \quad (77a)$$

$$Y = (1 - \frac{1}{2}\alpha \sin \omega t) x^2 \quad (77b)$$

$$(77c)$$

then

$$\dot{X} = \frac{dX}{dt} = \frac{1}{2}\alpha\omega \cos \omega t x^1 \quad (78a)$$

$$\dot{Y} = \frac{dY}{dt} = -\frac{1}{2}\alpha\omega \cos \omega t x^2 \quad (78b)$$

A plane gravitational wave of the form (72) with $f[(x^0 - x^3), y^3] = \alpha \sin[\omega(x^0 - x^3)] y^3$, and $y^3 = (0, 0, 1, 0)$ propagates in the y^3 direction. Some test masses in the $x - y$ plane are at rest in a circle about a central test mass. After the gravitational wave passes in time t the circle is squeezed in the Y -direction and expanded in the X -direction, therefore the circle is transformed to an elliptic shape.

We define the indicatrix curve I_p to be a circle with center 0 and radius $\sqrt{1 + f(x, y)}$ OP where $P(x, y)$ is an arbitrary point and $f(x, y)$ is a positive valued function. We can apply the Randers metric in order to calculate the distances between the masses [15]. Therefore we have

$$\mathcal{L}(x, y) = \frac{\lambda + \sqrt{f(x, y)\rho^2 + \lambda^2}}{f(x, y)} \quad (79)$$

where

$$\rho^2 = \frac{\dot{X}^2 + \dot{Y}^2}{X^2 + Y^2} \quad \lambda = \frac{X\dot{X} + Y\dot{Y}}{X^2 + Y^2} \quad (80)$$

Conclusions

We studied the behaviour of particles moving in a gravitational and electromagnetic field with the physical geometry of a Finsler–Randers (FR) space. Cartan and Berwald connections are applied for studying a linearized version of a weak field limit in F-R spaces.

In virtue of curvature tensors of the space of considerations some physical characterizations and interpretations in the sense of a «gravito–electromagnetic curvature» are given. Such a concept could play a role in the bending of light geodesics and gravitational lensing in a region of locally anisotropic space-time.

In paragraph 3 the Raychaudhuri equations are extended and they were derived in the framework of a tangent bundle. This consideration can give an additional interest to a string theory.

Finally, some applications of Randers metric for gravitational waves are presented.

Acknowledgement

The author is grateful to the University of Athens (Special Accounts for Research Grants) for the support to this work.

References

- [1] Weber J. : *Detection and Generation of Gravitational Waves* Phys.Review Vol.117, No1 (1960).
- [2] Wagoner R. : *Scalar Tensor and Gravitational Waves* Phys.Review D Vol. 1, No12 (1970).
- [3] K.Thorne and S. Kovacs : "The Generation of Gravitational Waves. I.Weak-Field Sources." The Astroph. Journal, 200: pp245-262, (1975)
- [4] Farese Es.: *Comparing Solar-System, Binary - Pulsar and Gravitational - Waves Tests of Gravity.* arxiv: gr-qc/9903058v1
- [5] B.Sathyaprakash, B.Shutz : *Physics,Astrophysics and Cosmology with Gravitational Waves*,Living Rev. Relativity 12 (2009),2.
- [6] Stavrinou, P.: *Deviations of Geodesics and Gravitational waves in Finsler space-time*, Rev. Bulletin of Calcutta Mathematical Society 9, pp. 1-12.
- [7] Stavrinou P.: *Gravitational and cosmological considerations based on the Finsler and Lagrange metric structures*, Nonlinear Analysis (2009), doi:10.1016/j.na.2009.01.182
- [8] P.C.Stavrinou : *On the linearized field theory of Finsler and Lagrange Spaces* Algebras Groups and Geometry 17 (2000),351-360
- [9] V.Balan and P.C.Stavrinou : *Weak Gravitational Fields in Generalized Metric Spaces* Geometry Balkan Press Proc.6 27-37 (2001)
- [10] N.Brinzei and S.Siparou : *On the possibility of the OMPR effect in the space with Finsler geometry I* pp41 Hpercomplex numbers in geometry and physics 2(8) Tom 4 (2007)
- [11] Randers, G., : *On an Asymmetrical Metric in the Four-Space of General Relativity*, Phys.Rev. Jan.15,(1941), p.191.
- [12] Bao,D. S-S Chern and Z.Schen: *An introduction to Riemannian-Finsler Geometry*, Springer New York (2000)

- [13] Stavrinou, P., Kouretsis, A., Stathakopoulos, M.: *Friedman-like Robertson-Walker model in generalized metric space-time with weak anisotropy*, General Relativity and Gravitation **40**, (2007), pp. 1403-1425.
- [14] Z.Chang,M-H Li,Xin Li,S.Wang : *Modified Rindler Potential in Randers-Finslerian space-time and the Convergence V-Map of Bullet Cluster* 1EO657558 arXiv : 11103893 v1 18-10-2011
- [15] Antonelli, P.L., Ingarden, R.S., Matsumoto, M.: *The Theory of Sprays and Finsler Spaces with applications in Physics and Biology*, Kluwer Academic Publishers, (1993).
- [16] Pfeifer C. and Wohlfarth M. arXiv : 1104.1079 v1.6 Ap.2011 Phys.Rev. D. 84044039
- [17] A.Bejancu and H.R. Farran : *Geometry of Pseudo-Finsler Submanifolds* Klumer Academic,Dordrecht(2000)
- [18] L.Iorio,C.Corda : *Gravitomagnetism and gravitational waves* arxiv : 1001.3951V2
- [19] Martens, R.:*Nonlinear gravito - electromagnetism*. General Relativity and Gravitation **40**, (2008), pp. 1203-1217.
- [20] Mashoon, B.:*Gravitomagnetism: A Brief Introduction*. arXiv: gr-qc/0311030.
- [21] Bel, L.: C. R. Acad. Sci. **247**, (1958), pp 1094.
- [22] Bel, L.: General Relativity and Gravitation **32**, (2000), pp 2047 (translation of 1962 article).
- [23] Matte, A.: Can.J. Math. **5**, (1953), pp 1.
- [24] Penrose, R.: Ann Phys, **10**, (1960), pp 171.
- [25] Pirani, F.: Phys. Rev. **105**, (1957), pp 1089.
- [26] Tsagas C.:*Gravitational Waves and Cosmic Magnetism*. Class.Quantum.Grav.19 (2007)pp.3709-3722.

- [27] Stavrinou, P., Diakogiannis, F. : *Finslerian Structure of Anisotropic Gravitational Field* Gravitation and Cosmology V.10,N.4(40) pp1-11 (2004) Russian Gravitational Society
- [28] Vacaru S.,Stavrinou P.,Gabourov E.,Gontsa D. : *Clifford and Riemann-Finsler Structures in Geometrical Mechanics and Gravity* Geometry Balkan Press pp.643 (2005)
- [29] Vacaru S. : *Super-Luminal Effects for Finsler Branes as a way to preserve the Paradigm of Relativity Theories* arXiv :1110.0675v1 3 Oct 2011
- [30] A.Kostelecky : *Riemann-Finsler geometry and Lorentz-violating kinematics* Phys. Lett B701 (2011) pp 137-143
- [31] Asanov, G.S.: *Finsler Geometry, Relativity and Gauge Theories*, Reidel Pub. Comp. (1985).
- [32] Yasuda, H., Shimada, H.: *On Randers space of scalar curvature* Rep. Math. Phys. **11**, (1977), pp 347-360.
- [33] GS.Asanov and P.C.Stavrinou : *Finslerian deviations of geodesics over tangent bundle* Rep. Moth Physics (1991) pp 63-69
- [34] P.Stavrinou : *Tidal Forces in Vertical Spaces of Finslerian Space-Time* Rep on Math.Phys.(1992)Vol.31 pp1-4
- [35] S.Rutz : *Symmetry in Finsler Spaces*, Finsler Geometry Research Conf on Finsler Geometry July 16-20 1995 Seattle Eds D.Bao, S-S Chern and Z.Shen, Contemporary Mathematics pp 289
- [36] V.Balan and Stavrinou P. : *Deviations of Geodesics in Fibered Finslerian Approach* P.L.Antonelli and R.Miron, Lagrange and Finsler Geometry pp.65-74 Kluwer Acad.Pub (1996)
- [37] R.Miron and M.Anastasiei : *The Geometry of Lagrange Theory and Applications* Klumer Academic Dordrecht(2000)
- [38] Raychaudhuri.A : *Relativistic Cosmology* Phys.Rev.Vol.98 Numb 4 pp.1123,(1995)
- [39] S.Hawking-Ellis : *The large Scale Structure of Space-Time* Cambridge University Press (1973)

- [40] A.Kouretsis and C.Tsagas : *Raychaudhuri's equation and aspects of relativistic Charged collapse*"Phys.Rev.D82(2010) 124053
- [41] C.M.Claudel, K.S. Virbhadra, G.F.R. Ellis : *The geometry of photon surfaces* Jour. of Math. Physics. Vol.42, Numb. 2, Feb(2001)
- [42] K.S.Virbhadra and G.F.R. Ellis : *Schwarzschild black hole lensing* Phys. Rev. D. Vol. 62, 084003-1(2000)
- [43] P.C.Stavrinos : *Congruences of Fluids in a Finslerian Anisotropic Space-Time* Int Journ of Theor.Physics Vol44 No2 Feb(2005) 245-254